


Initiated: 9/92
Reviewed/revised: 2/13/08
Revision: 5

**MILWAUKEE COUNTY EMS  
MEDICAL PROTOCOL  
SEIZURE**

Approved by: Ronald Pirrallo, MD, MHSA
Signature:
Page 1 of 1

<b>History:</b>	<b>Signs/Symptoms:</b>	<b>Working Assessment:</b>
Reported/witnessed seizure activity History of seizures Medic alert tag Anti-seizure medications History of recent trauma History of diabetes Pregnancy Fever	Seizure activity Decreased mental status (post ictal) Sleepiness Incontinence Trauma	Seizure (look for underlying cause): <ul style="list-style-type: none"> <li>• Head trauma</li> <li>• Noncompliance</li> <li>• Fever/infection</li> <li>• Hypoglycemia</li> <li>• Overdose/poisoning</li> <li>• Alcohol withdrawal</li> <li>• Hypoxia</li> <li>• Eclampsia</li> </ul>



**NOTE:**

- Pediatric patients with febrile seizures rarely seize more than once. If patient seizes again, evaluate for another cause.
- Status Epilepticus is defined as two or more successive seizures without a period of consciousness or recovery.
- AEIOU-TIPS-V = A - alcohol, airway, arrest; E - epilepsy, electrolytes, endocrine; I - insulin; O - overdose, oxygen depletion, opiates; U - Uremia/chronic organ failure; T - trauma, tumors, temperature; I - infection; P - psychiatric, pseudoseizures; S - Syncope, shock, stroke, sickle cell crisis; V - vascular/lack of blood flow.